

Nurses' Knowledge Regarding Prevention of Ventilator Associated Pneumonia

Aurang Zeb*, Muhammad Hasnain, Ijaz Ahmad, Shafiq Khan and Aftab Ali shah

Assistant Professor Rehman College of Nursing Peshawar, Pakistan

Received: 📅 November 16, 2018; Published: 📅 November 27, 2018

*Corresponding author: Aurang Zeb, Assistant Professor Rehman College of Nursing Peshawar, Pakistan

Abstract

Ventilator Associated Pneumonia (VAP) is one of the common nosocomial infections with the prevalence rate of 10% to 70% in Intensive Care Units (ICUs). The incidence of VAP is 22.8% in patients on mechanical ventilation. Nurse's lack of knowledge about infection prevention and proper nursing care may become barrier to Prevent VAP.

Aim: The Aim of this study is to explore nurse's knowledge regarding Prevention of VAP.

Methods: Cross-sectional-study was conducted at both public and private tertiary care hospitals in Peshawar from March to April 2018. Nurse's knowledge was assessed by using close-ended questionnaire from 100 Critical-care Nurses. Participants were selected through convenient sampling. The questionnaire was adopted from similar studies which included 15 questions. Data was analyzed using descriptive statistics by SPSS software version 22.

Results: This study consisted of 100 Participants including 30% males and 70% females. 33% of the participants were married and 67% were unmarried. Professional qualification of participants was also identified, 53% of the participants were diploma holders and 47% were generic BSN. The mean ICU experiences of participants were 2.79+3.0 years. The mean Score in the knowledge part of VAP prevention out of 15 marks was 8.14+ 1.8.

Conclusion: The study concludes that several educational programs shall be arranged for nurses to improve their knowledge of VAP prevention and can decrease Ventilator associated Pneumonia among patients.

Introduction

Pneumonia is described as an inflammation of lung that primarily affects alveoli McLuckie [1]; Leach & Richard. Pneumonia is usually caused by the infection with virus, bacteria and another microorganism McLuckie [1]; Jeffrey & Pommerville [2]. Ventilator associated pneumonia (VAP) is one of the common nosocomial infections with the prevalence rate of 10% to 70% in critical care units Ruffell [3]. Ventilator-associated pneumonia is a type of lungs infection that occurs in people on mechanical ventilator in hospitals. It typically affects source of increased illness and death. Patients with ventilator acquired pneumonia (VAP) have increased Stay-length in hospital and have up to a 20-30% death rate Cook [4]. The incidence of VAP is 22.8% in patients on mechanical ventilation in developed countries Safdar N; Dezfulian C [5]. Ventilator associated Pneumonia is mainly caused after aspiration when the pathogen is transmitted to critically ill Patients in intensive care unit Michetti [6].

Aspiration causes Oro-pharyngeal Colonization and infection. Most of the patients admitted in ICU are already colonized with

bacterial infection Ruffel [3]. Ventilator-associated Pneumonia (VAP) is the common nosocomial infection which mostly occurs in ICU's Patients on Mechanical Ventilation for 48 hours or longer; it increases hospital stay and financial cost of the Patients Cooper & Haut [7]; Yazdani [8]. Furthermore, VAP is also associated with prolonged ICU stay in hospital and delayed extubating, that increases morbidity and mortality rate El-Khatib [9]. The study "Critical Care Nurse's Knowledge of Ventilator-Associated Pneumonia Prevention in a Hospital in Khartoum" by Al Shameri [10] resulted that nurses have inadequate knowledge about VAP prevention.

Another study "Critical Care Nurses' Knowledge and Compliance with Ventilator Associated Pneumonia Bundle at Cairo University Hospitals" in Cairo, Egypt by Nahla [11] indicates that all critical care nurses, with different educational levels, have unsatisfactory knowledge scores about ventilator associated pneumonia and VAP bundle preventive measure. A cross-sectional study "Knowledge Levels of Intensive Care Nurses on Prevention of Ventilator-associated Pneumonia" in Turkey by Korhan [12] shows that critical care nurses have poor knowledge about ventilator

associated Pneumonia. A descriptive study “Nurses’ knowledge of evidence- based guidelines for preventing ventilator associated pneumonia in intensive care units” in Iran by Masoumeh Bagheri-Nesam [13] quantified the result in percentages and showed that level of the nurses’ knowledge about preventive measures was 51.92%; they also recommended the need of more educational programs in the target field. A study to assess the knowledge and practice of intensive care nurses on prevention of ventilator-associated pneumonia (VAP) among patients admitted in critical care units of Government Medical College Hospitals, Jammu (J&K), by Jyoti Kapoor [14] revealed that ICU nurses had average knowledge regarding ventilator acquired Pneumonia. A study on the same topic in Karachi, Pakistan by Yasmin [15] resulted that nurse working at critical unit were having knowledge-gap to be able to prevent VAP among ventilated patients.

Nurses’ Role in Prevention of VAP

Critical care nurses play an important role in prevention of Ventilator associated Pneumonia by preventing patient from risk factors, notifying early symptoms of VAP in Patients and assisting in diagnosis Myrianthefs [16]. Nurses are health care providers who protect patients from infection. So, lack of knowledge about the prevention of infection and proper nursing care may become barrier to prevent VAP. The occurrence of ventilator acquired Pneumonia indicates inadequacy of experienced nurses, insufficient knowledge and understanding about the pathophysiology and prevention of VAP Augustyn [17]; Augustyn.

The spread of VAP may be due to ICU Nurses’ contact with patients receiving care of mechanical ventilation and their contact with nursing students, visitors and health care workers. Moreover, they are responsible for delivering nursing care to patient. As nurses deliver care related to mechanical ventilation, they should have appropriate knowledge regarding ventilator moods, positive pressure ventilation, daily weaning process, aspiration prevention and head on the bed elevation. This knowledge should be integrated into Practice to Prevent VAP Keyt, Faverio, & Res [18]; Gonçalves [19].

However, ventilator associated pneumonia is preventable by educating ICUs nurses regarding VAP. No study has been found on knowledge of intensive care nurses regarding prevention of VAP in KPK. Therefore, it is important to explore intensive care nurse’s knowledge, and to highlight deficiency in knowledge for creating awareness and methods to prevent VAP in KPK which is the aim of this study.

Purpose of the Study

The purpose of this study is to explore nurses’ knowledge regarding Prevention of VAP in order to prevent complications.

Methodology

Study Design: Descriptive cross-sectional was conducted at

both public and private tertiary care hospitals in Peshawar from March to April 2018.

Population and Setting: Both public and private sector hospitals were selected for conducting this study. The population of this study was ICU Nurses in hospital.

Sampling Technique: Critical Care Nurses of different ICUs were selected by using Convenient Sampling technique.

Sample Size: Sample size is calculated by Raosoft software, with 5% margin of error and 10% non-response rate the proportionate sample obtained is given below. So, the total sample size from three hospitals is 100.

Inclusion Criteria: Registered nurses working in ICUs.

Exclusion Criteria: ICU nurses with less than six months of experience. Managerial nurses do not directly involve in patient care.


Data Collection Tool: An adopted well organized questionnaire was used for data collection Ally Tatu Said;Labeau. The questionnaire contained fifteen questions about Prevention of Ventilator-acquired Pneumonia.



Ethical Consideration: Approval for the data collection was taken from the Hospital Directors. Written informed consent was presented to each participant for his/her agreement as a participant. Anonymity was guaranteed to all the participants. The whole data was kept confidential. Only primary data collectors and supervisors had access [20-23].

Data Analysis

Data was analyzed by using SPSS version 22. In descriptive statistics, frequencies and percentages were calculated for nominal and ordinal data while mean and standard deviation were calculated for continuous variables. In inferential statistics paired T test was applied to identify the effectiveness of informed consent on preoperative anxiety.

Result





This study consisted of 100 participants including 30% males and 70% females (Chart 1 & 2). 33% of the participants were married and 67% were unmarried (Table 1). As for as Academic qualification of the participants is concerned, 30% participants had Matric, and 70% participants had FA/FSC qualification (Chart 3).

Table 1:

	Frequency	Percent
Married	33	33
Unmarried	67	67
Total	100	100

Discussion

Literature is available on the subject matter; studies have been conducted in different parts of the world. These studies show different results [24]. Some of them are related to the result of current study (Chart 4). A study “Critical Care Nurse’s Knowledge of Ventilator-Associated Pneumonia Prevention was done in Hospitals in Khartoum” by Al Shameri [10] in which data were collected through the use of structured knowledge questionnaire filled from 120 ICU nurses (Table 2). The result showed that the overall knowledge of nurses regarding the VAP prevention mean was (18.22 from 40) that was a poor result. This shows us that most

of the nurses have inadequate knowledge about VAP prevention (Table 3).




Table 2:

	Frequency	Percent
Diploma	53	53
BSN	47	47
Total	100	100

Table 3:

	N	Mean	Std. Deviation
Total experience	100	4.27	4.5
ICU experience	100	2.79	3
Score of participants	100	8.14	1.8

Another study “Critical Care Nurses’ Knowledge and Compliance with Ventilator Associated Pneumonia Bundle at Cairo University Hospitals” in Cairo, Egypt was conducted by NahlaShaaban Ali [11] in which data were collected from 45 ICU nurses through Self-administered questionnaire. The result showed that all critical care nurses with different educational levels, had unsatisfactory knowledge scores about ventilator associated pneumonia and VAP bundle preventive measure [25]. In cross-sectional study titled as “knowledge levels of intensive care nurses on prevention of ventilator-associated Pneumonia” in Turkey, Korhan [12] collected data by means of a Nurse Identification Form and a Form of Evidence-Based Knowledge regarding the Prevention of Ventilator-Associated pneumonia; in that study, the findings were with the median value of total points scored by nurses on the questionnaire was 4.00 ± 2.00. And the median value of their total scores on the questionnaire was found to be statistically significant (p < 0.05). The conclusion of the study was that critical care nurses’ knowledge about ventilator-associated pneumonia prevention was poor Korhan [12].

In a descriptive study “Nurses’ knowledge of evidence- based guidelines for preventing ventilator associated pneumonia in

intensive care units” in Iran, Bagheri-Nesami and Amiri [13] assessed the knowledge of 52 ICU nurses through questionnaire, and they quantified the result in percentages which showed that showed the level of the nurses’ knowledge about preventive measures was 51.92%. They also recommended the need of more educational programs in this field. Similarly, Jyoti Kapoor (2017) found that ‘intensive care nurses have average knowledge and unsatisfactory practice regarding prevention of ventilator-associated pneumonia.’ While, the current study shows that among 100 ICU nurses, 88% nurses have average knowledge, 5% of them has excellent knowledge and 7% nurses have poor knowledge about ventilator acquired pneumonia prevention.

Conclusion and Recommendations

Current study revealed that 80% nurses of ICU have average knowledge; while, only 5% nurses have excellent knowledge regarding VAP. Thus, it is concluded that several educational programs should be arranged for nurses to improve their knowledge from average to excellent and to decrease Ventilator associated Pneumonia among patients.

References

1. Leach Richard E (2009) *Acute and Critical Care Medicine at a Glance* (2nd ed.). Wiley-Blackwell. McLuckie.
2. Jeffrey C and Pommerville (2010) *Alcarno's Fundamentals of Microbiology* (9th ed.). Sudbury MA: Jones & Bartlett. pp. 323.
3. Ruffell A and Adamcova L (2008) Ventilator-associated pneumonia: prevention is better than cure. *Nurs Crit Care* 13(1): 44-53.
4. Cook D (2000) Ventilator associated pneumonia: perspectives on the burden of illness. *Intensive Care Med* 26(1): 31-37.
5. Safdar N, Dezfulian C, Collard H R, Saint S (2005) Clinical and economic consequences of ventilator-associated pneumonia: A systematic review. *Crit Care Med* 33(10): 2184-2193.
6. Michetti C P, Fakhry S M, Ferguson P L, Cook A, Moore FO, et.al (2012) Ventilator-associated pneumonia rates at major trauma centers compared with a national benchmark: a multi-institutional study of the AAST. *J Trauma Acute Care Surg* 72(5): 1165-1173.
7. Cooper V B and Haut C (2013) Preventing Ventilator-Associated Pneumonia in Children: An Evidence-Based Protocol. *Critical Care Nurse* 33(3): 21-29.
8. Yazdani M, Sabetian G, Ofish R A, Roudgari A, Feizi M (2015) The effect of nurses training on ventilator-associated pneumonia (VAP) prevention bundle on VAP incidence rate at a critical care unit. *Journal of Nursing Education and Practice* 5(12).
9. El Khatib, M F, Zeineldine S, Ayoub C, Husari A, Bou Khalil P K (2009) Critical Care Clinicians’ Knowledge of Evidence-Based Guidelines for Preventing Ventilator-Associated Pneumonia. *Am J Crit Care* 19(3): 272-276.
10. Al Shameri FA (2017) *Critical Care Nurse’s Knowledge of Ventilator-Associated Pneumonia Prevention Selected Hospitals, Khartoum*.
11. Nahla Shaaban Ali (2013) *Critical Care Nurses’ Knowledge and Compliance with Ventilator Associated Pneumonia Bundle at Cairo University Hospitals*. *Journal of Education and Practice* 4(15).
12. Akin Korhan E, HakverdioğluYönt G, Parlar Kılıç S, Uzelli D (2013) Knowledge levels of intensive care nurses on prevention of ventilator associated pneumonia. *Nurs Crit Care* 19(1): 26-33.
13. Bagheri Nesami M and Amiri M (2014) Nurses’ knowledge of evidence-based guidelines for preventing ventilator-associated pneumonia in intensive care units. *Journal of Nursing and Midwifery Sciences* 1(1): 44-48.
14. Kapoor J (2017) knowledge and practice of intensive care nurses on prevention of ventilator-associated pneumonia. *VAP National Journal of Multidisciplinary Research and Development* 2(3): 358-360.
15. Meherali S, Parpio Y, Ali T S, Javed F (2011) Nurses’ knowledge of evidence-based guidelines for prevention of ventilator associated pneumonia in critical care areas. pre and post testdesign. *J Ayub Med Coll Abbottabad* 23(1): 146-149.
16. Myrianthefs P M, Kalafati M, Samara I, Baltopoulos G J (2004) Nosocomial Pneumonia. *Critical Care Nursing Quarterly* 27(3): 241-257.
17. Augustyn B (2007) Ventilator-Associated Pneumonia: Risk Factor and Prevention. *Crit Care Nurse* 27(4): 32-36.
18. Keyt H, Faverio P, Restrepo M I (2014) Prevention of ventilator associated pneumonia in the intensive care unit. a review of the clinically relevant recent advancements. *Indian J Med Res* 139(6): 814-21.
19. Gonçalves F A, Brasil V V, Ribeiro L C, Tipple A F (2012) Nursing actions for the prevention of ventilator-associated pneumonia. *Acta Paul. Enferm* 25: 101-107.
20. Dodek P, Keenan S, Cook D, Heyland D, Jacka M, et.al (2004) Evidence-Based Clinical Practice Guideline for the Prevention of Ventilator-Associated Pneumonia. *Ann Intern Med* 141(4): 305-313.
21. Labeau S, Vandijck D, P Van Aken, S Blot, Claes B (2007) Intensive care nurses’ knowledge of evidence-based guidelines for the prevention of ventilator-associated pneumonia. *Crit Care* 11(2): 92.
22. Mc Luckie A (2009) Pneumonia. *Respiratory Disease and its Management* pp. 51-59.
23. Resende M M, Monteiro S G, Callegari B, Figueiredo P M, Monteiro C R, et.al (2013) Epidemiology and outcomes of ventilator-associated pneumonia in northern Brazil: an analytical descriptive prospective cohort study. *BMC Infect Dis* 13(1): 119.
24. Tablan O C, Anderson L J, Besser R, Bridges C, Hajjeh R (2003) Guidelines for Preventing Health-Care-Associated Pneumonia, 2003: Recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee. *MMWR Recomm Rep* 53(3): 1-36.
25. Zack J E, Garrison T, Trovillion E, Clinkscale D, Coopersmith C M, et.al (2002) Effect of an education program aimed at reducing the occurrence of ventilator-associated pneumonia. *Crit Care Med* 30(11): 2407-2412.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Article](#)

DOI: [10.32474/LOJNHC.2018.02.000129](https://doi.org/10.32474/LOJNHC.2018.02.000129)



Lupine Online Journal of Nursing & Health care

Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles